



TrichoConcept™

Compatibility Table



Table 1.
List of compatible APIs/DCIs



API/DCI	Range of concentration (%)	TrichoSol™	TrichoFoam™	TrichoSerum™	TrichoCream™	TrichoOil™	TrichoWash™	TrichoCond™
17- α -Estradiol	0.025 to 0.05	● / ●	● / ●		*			
Adenosine	0.5 to 0.75	*					*	
Arginine HCl	0.5 to 2.0			*			*	*
Betamethasone dipropionate	0.05	*	*					
Betamethasone dipropionate	0.025 to 0.05				*			
Betamethasone valerate	0.1	*	*					
Biotin	0.5 to 2.0	*						
Biotin	0.1 to 0.4			*			*	*
CafeiSome™	1.0 to 5.0	*	*					
CafeiSome™	2.0 to 5.0						*	
Caffeine	0.1 to 2.0	●	●				*	
Cetirizine HCl	0.5 to 2.0	●	*					
Ciclopirox olamine	1.5						*	
Clobetasol propionate	0.01 to 0.1	●	● / ●					
Clobetasol propionate	0.05				*		●	
Clotrimazole	1.0					**	*	
Coal tar	1.0 to 1.5						*	
Cyproterone acetate	1.0	*	*					
Desonide	0.025 to 0.05	*	*		*			
D-panthenol	0.25 to 0.75		*		*			
D-panthenol	0.5 to 5.0			*			*	*
Dutasteride	0.1 to 0.25	● / ●	● / ●					
Erythromycin	0.5 to 4.0					**		
Erythromycin	2.0 to 4.0						*	
Estrone	0.01 to 0.05	*						
Finasteride	0.1	*	●					
Finasteride	0.1 to 0.25	●	●					
Finasteride	0.5 to 1.0				●			
Fluocinolone acetonide	0.01	*	*		*	**	*	
Ginkgo biloba glycolic extract	1.5 to 3.0	*	*	*	*	**	*	*
Ginseng glycolic extract	1.5 to 3.0	*	*	*	*	**	*	*
Hydrocortisone acetate	0.5 to 1.0	● / ●						
Hydrocortisone acetate	1.0				*			

Compatible combination for

* 35 days EBUD ** 180 days EBUD ● 7 days ● 14 days ● 30 days ● 60 days ● 90 days ● 120 days ● 150 days ● 180 days ● 380 days ○ not compatible

* Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 35 days for preserved aqueous formulations.
** Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 180 days for non-aqueous formulations.





API/DCI	Range of concentration (%)	TrichoSol™	TrichoFoam™	TrichoSerum™	TrichoCream™	TrichoOil™	TrichoWash™	TrichoCond™
Hydrocortisone butyrate	0.1	*	*					
IGrantine-F1™	0.1 to 5.0	*	*		*			
Ketoconazole	2.0	*	*				●	
Latanoprost	0.005	*	*		*			
Latanoprost	0.05	●	●					
Latanoprost	0.1				●			
L-Carnitine L-tartrate	0.5 to 2.0	*	*				*	
Melatonin	0.0033 to 0.1	*	*					
Metformin HCl	5.0 to 10.0	*						
Metronidazole	0.75	●			*			
Minoxidil	1.0 to 7.0	● / ●	● / ●					
Minoxidil	2.0 to 5.0				*			*
Nicotinamide	0.25 to 0.5	●	●	*		**	*	*
Prednicarbate	0.05 to 0.1					**		
Prednicarbate	0.1		●					
Progesterone	0.25 to 2.5		● / ●					
Progesterone	1.0 to 2.5	● / ●						
Prostaquinon	1.5 to 3.0	*	*		*			
Prostaquinon	3.0					●		
Pyridoxine HCl	0.25 to 5.0	●						
Pyridoxine HCl	0.5			*			*	*
Retinol palmitate	0.5 to 1.0			*		**	*	*
Salicylic acid	6.0					**	*	
Salicylic acid	2.0 to 6.0				*		*	*
Serenoa serrulata dry extract	1.0 to 2.0	*	*					
Serenoa serrulata lipid extract	1.0 to 2.0				*	**	*	*
SiliciuMax™	5.0 to 10.0	*	*	*	*		*	*
Spironolactone	1.0	●	●				●	
Tacrolimus	0.1	*				**		
Tocopherol acetate	1.0 to 2.0			*		**	*	*
Tretinoin	1.0	*	*					
Triamcinolone acetonide	0.1	*	*					
TrichoXidi™	2.5	●	*					
Urea	5.0 to 10.0					**	*	
VEGF TrichoFactor™	2.5	*	*					
Zinc pyrithione	2.0						*	

Compatible combination for

* 35 days EBUD ** 180 days EBUD ● 7 days ● 14 days ● 30 days ● 60 days ● 90 days ● 120 days ● 150 days ● 180 days ● 380 days ○ not compatible

* Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 35 days for preserved aqueous formulations.

** Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 180 days for non-aqueous formulations.



Table 2.
List of compatible
APIs/DCIs associations



API/DCI association	Range of concentration (%)	TrichoSol™	TrichoFoam™	TrichoSerum™	TrichoCream™	TrichoOil™	TrichoWash™	TrichoCond™
Caffeine + Pyridoxine + Menthol	2.0 + 0.5 + 1.0						●	
Latanoprost + 17- α -estradiol	0.005 + 0.05	*	*					
Latanoprost + Dutasteride + Ginkgo biloba glycolic extract	0.005 + 0.25 + 2.25	*	*					
Latanoprost + Spironolactone	0.005 + 0.52	*						
Minoxidil + 17- α -estradiol	5.0 + 0.05	*						
Minoxidil + 17- α -estradiol + Arginine HCl	2.0 + 0.05 + 0.25	*						
Minoxidil + Betamethasone valerate	5.0 + 0.1	*						
Minoxidil + Clobetasol	5.0 + 0.05	*						
Minoxidil + Dutasteride	5.0 + 0.1 / 7.0 + 0.25	* / ●	* / ●					
Minoxidil + Finasteride	1.0 + 0.1 / 7.0 + 0.25	* / ●	● / ●					
Minoxidil + Finasteride + Tretinoin		⊗						
Minoxidil + Hydrocortisone butyrate	5.0 + 0.1	*						
Minoxidil + Ketoconazole	1.0 + 0.5 / 7.0 + 2.0	●	● / ●					
Minoxidil + Latanoprost	5.0 + 0.005	*	*					
Minoxidil + Latanoprost + Finasteride	5.0 + 0.1 + 0.25	*	*					
Minoxidil + Tretinoin		⊗						
Minoxidil + Triamcinolone acetonide	5.0 + 0.2	*						
Minoxidil + TrichoXidi™	2.5 + 2.5	●						
Prednisolone + Clotrimazole		⊗						

Compatible combination for

* 35 days EBUD ** 180 days EBUD ● 7 days ● 14 days ● 30 days ● 60 days ● 90 days ● 120 days ● 150 days ● 180 days ● 380 days ⊗ not compatible

* Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 35 days for preserved aqueous formulations.
** Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 180 days for non-aqueous formulations.





API/DCI association	Range of concentration (%)	TrichoSol™	TrichoFoam™	TrichoSerum™	TrichoCream™	TrichoOil™	TrichoWash™	TrichoCond™
Pyridoxine + Retinol palmitate + Tocopherol acetate	0.5 + 1.0 + 12.1IU/g							●
Spironolactone + Progesterone + Minoxidil		⊗						
Sulfur + Salicylic acid		⊗						
Triamcinolone + Clotrimazole			⊗					

Compatible combination for

* 35 days EBUD ** 180 days EBUD ● 7 days ● 14 days ● 30 days ● 60 days ● 90 days ● 120 days ● 150 days ● 180 days ● 380 days ⊗ not compatible

* Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 35 days for preserved aqueous formulations.
** Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 180 days for non-aqueous formulations.





Disclaimer

While a great deal of effort has been made to ensure the accuracy of the information contained herein, no claims are made as to the uses, safety, efficacy, or bioavailability of these recommendations. The content of this document cannot be construed as (medical) advice, recommendation, or opinion. Health professionals using this information are advised to do so solely if appropriate in their own professional opinion and judgment. Fagron does not accept and cannot be held responsible or liable in any case for the information contained herein.

Fagron BV

Fascinatio Boulevard 350
3065 WB Rotterdam
The Netherlands

T +31 88 33 11 288
F +31 88 33 11 210
www.fagron.com